

8400011

TO ALL TO WHOM THESE: PRESENTS: SHALL COME;

Minnesota Agricultural Experiment Station Colherens, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT


UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF Eighteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR JUSTIC IT IN PRODUCING A HYBRID OR DIFFERENT ETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT.

UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

BARLEY

'Robust'

In Testimony Whereof, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington this 29th day of March the year of our Lord one thousand nine hundred and eighty-five.



U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, MEAT, GRAIN & SEED DIVISION

APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE

FORM APPROVED: OMB NO. 0581-0055
Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).

(Instructions on reverse		ION CENTIFICATE		confidential until certificate is issued S.C. 2426).
1. NAME OF APPLICANT(S)	2.	TEMPORARY DESIGNATION	3. V	ARIETY NAME
Minnesota Agricultural Experiment Statio	on	Minn. M36	R	obust
4. ADDRESS (Street and No. or R.F.D. No., City, State, and Zip C		PHONE (Include area code)		FOR OFFICIAL USE ONLY
Univ. of MN, 220 Coffey Hall			PVPC	NUMBER
1420 Eckles Ave., St. Paul, MN 55108		(612) 373-0751	J 44	8400011
6. GENUS AND SPECIES NAME 7. FAMILY	YNAME	(Botanical)	(2)	DATE
Hordeum vulgare L. Gram	inae		FILING	11/1/83 TIME 2:30 A.M. XP.M.
8. KIND NAME	9. D	ATE OF DETERMINATION		AMOUNT FOR FILING
Barley, six-rowed	F	eb. 15, 1983	RECEIVED	DATE 11/1/83
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE For partnership, association, etc.)	ORM OF	ORGANIZATION (Corporation,	FEES RE	s 500.00 DATE 3-28-85
11. IF INCORPORATED, GIVE STATE OF INCORPORATION			12. [DATE OF INCORPORATION
Department of Agronomy and Plant Gene University of Minnesota St. Paul, MN 55108 14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SU. a. Exhibit A, Origin and Breeding History of the Variety (See Section 52 of the Plant Variety Protection Act.) b. Exhibit B, Novelty Statement 15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS SEED? (See Section 83(a) of the Plant Variety Protection Act.)	UBMITTE See VARIET	c. X Exhibit C, Objective D from Plant Variety Productional	Descri _j	Y AS A CLASS OF CERTIFIED
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY LIMITED AS TO NUMBER OF GENERATIONS?	/ BE	17. IF "YES" TO ITEM 16, V BEYOND BREEDER SEE	VHICH	
X Yes No		X Foundation	_	egistered X Certified
18. DID THE APPLICANT(S) FILE FOR PROTECTION OF THE	VARIE	TY IN THE U.S.?		Yes (If "Yes," give date) No
19. HAS THE VARIETY BEEN OFFERED FOR SALE OR MARK				Yes (If "Yes," give names
The variety has been released and is 1983 and 1984.	s beir	ig marketed in the U	.5.	of countries and dates)
20. The applicant(s) declare(s) that a viable sample of basic plenished upon request in accordance with such regulat			with	the application and will be re-
The undersigned applicant(s) is (are) the owner(s) of the distinct, uniform, and stable as required in Section 41, a Variety Protection Act.	nis sexua and is e	ally reproduced novel plant var ntitled to protection under the	iety, a	and believe(s) that the variety is isions of Section 42 of the Plant
Applicant(s) is (are) informed that false representation	herein c	can jeopardize protection and		
SIGNATURE OF APPLICANT			D	ATE
a onald C Grasm	1 us	20n		Feb. 27, 1984
SIGNATURE OF APPLICANT			D	ATE

INSTRUCTIONS

General: Send an original copy of the application and exhibits, at least 2,500 viable seeds, and \$1,000 fee (\$500 filing fee and \$500 examination fee) to U.S. Department of Agriculture, Agricultural Marketing Service, Livestock, Meat, Grain and Seed Division, Plant Variety Protection Office, National Agricultural Library Building, Beltsville, Maryland 20705. (See section 180.175 of the Regulations and Rules of Practice.) Retain one copy for your files. All items on the face of the form are self-explanatory unless noted below.

Item

- Give the date the applicant determined that he had a new variety based on (1) the definition in section 41(a) of the Act and (2) the date a decision was made to increase the seed.
- Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method; (2) the details of subsequent stages of selection and multiplication; (3) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified and (4) evidence of uniformity and stability.
- Give a summary statement of the variety's novelty. Clearly state how this novel variety may be distinguished from all other varieties in the same crop. If the new variety most closely resembles one or a group of related varieties: (1) identify these varieties and state all differences objectively; (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and (3) submit, if helpful, seed and plant specimens or photographs of seed and plant comparisons clearly indicating novelty.
- 14c Fill in the Exhibit C, Objective Description form, for all characteristics for which you have adequate data.
- Describe any additional characteristics that are not described, or whose description cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the description of characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- If "Yes" is specified (seed of this variety be sold by variety name only as a class of certified seed) the applicant may NOT reverse his affirmative decision after the variety has either been sold and so labeled, his decision published, or the certificate has been issued. However, if the applicant specified "No," he may change his choice. (See section 180.16 of the Regulations and Rules of Practice.)
- See section 42 of the Plant Variety Protection Act and section 180.7 of the Regulations and Rules of Practice.

14A. Exhibit A, Origin and Breeding History of the Variety:

Robust originated from the cross 'Morex'/'Manker' made in 1973. The F₂ generation was grown in the field and individual plants were selected. The F3 and F4 generations were grown in the greenhouse using a single seed descent procedure. The F5 generation was grown in the field and selection was practiced for agronomic and malting quality traits. Robust traces to a single plant that was taken at random from a selected F5 line. Seed of the single plant was increased in a winter nursery in Mexico and replicated testing was begun in the F7 generation in 1976. Testing for disease and agronomic merit in statewide trials continued through 1982 and in a regional trial in 1979 through 1982. Malting and brewing quality was evaluated at the USDA Barley and Malt Laboratory, Madison, Wisconsin and in industry pilot and plant scale tests starting in 1975 (F5 generation) and continuing through 1982. Reselection was done in 1979 (Fig generation) when 45 head rows were bulked to reconstitute the line. Robust contains no off-type plants.

14B. Exhibit B, Novelty Statement:

Robust is a six-rowed, smooth-awned spring barley. The kernels are covered, medium size, with short rachilla hairs, and a white aleurone. The spike is medium-lax, medium-large, and semi-erect. It is medium late, mid-tall and has moderately strong straw.

Robust is intermediate to its parents, 'Morex' and 'Manker', in plant characters such as leaf size, head number and kernel number. It is, however, two to three days later in maturity than either parent. It is resistant to races of stem rust and spot blotch which occur in Minnesota. It is susceptible to prevalent races of loose smut.

Robust is different from all other midwestern malting barleys with which it might be compared. Of the midwestern barleys it is most similar to Morex, one of its parents. However, Robust is different from Morex in being 2 days later maturing, more resistant to lodging, higher yielding and in having a higher percentage of plump kernels (Table 1).

14D. Exhibit D, Additional Description of Robust

'Robust' barley (<u>Hordeum vulqare L.</u>) (Reg. No. 184), P.I. 476976, was developed by the Minnesota Agricultural Experiment Station and released 15 Feb. 1983. Robust originated from a single F5 plant selected from a 'Morex'/'Manker' cross made in 1973, and was tested as M36.

The breeding procedure was a combination of pedigree selection and single seed descent. Robust has been tested in all of the north central states and some testing has been done in the northwestern states. The proposed area for growing is the north central barley growing states. It may be suitable for growing in the western states, but this is not determined.





14D. (cont)

Robust is a six-rowed, smooth-awned spring barley. The kernels are covered and medium-large, with a short haired rachilla and white aleurone. The spike is medium-lax, medium-long, and semi-erect. Robust is medium-late, midtall, and has moderately strong straw.

Seed stocks will be maintained by the Minnesota Agricultural Experiment Station and the Minnesota Crop Improvement Association. Certification will be limited to three generations after breeder seed: Foundation, Registered and Certified. Certified seed will be offered for sale in 1983. Application will be made for protection via the "Certification Option".

14A. Exhibit A, addendum:

Robust is stable and can be maintained and reproduced by seed through a number of generations without changing its characteristics.



Table 1. Agronomic performance of Robust compared to Morex in Minnesota, 1979-83.

	No. of station years	Morex	Robust
Yield, bu/a	27	74	79
Plump barley, %	19	67	80
Lodging, %	19	36	28
Heading date, June	19	17	19



GALLEY1

9	000	CROP SCIENCE, VOL. 23, SEPTEMBER-OCTOBER 198
10		REGISTRATION OF CROP CULTIVARS
13		
14	ROBUST BA	RLEY
15	'ROBUST' barley (Hordeum vulgar	e L.) (Reg. No. 184), P.I.
16	476976, was developed by the M	innesota Agricultural Ex-
17	periment Station and released 1:	5 Feb. 1983. Robust orig-
18	inated from a single F5 plant s	
19	'Manker' cross made in 1973, an	
20	bust is a six-rowed, smooth-awne	
2i	nels are covered and medium-la	arge, with a short haired
22	rachilla and white aleurone. The	spike is medium-lax, me-
23	dium-long, and emicros. Robus	
24	and has moderately strong straw.	Although relatively widely
25	adapted, it is probably best suite	d for the barley growing
26	area of the upper midwest.	
27	Robust is superior to Morex, th	ne most widely grown cul-
28	tivar in Minnesota(1), in grain yie	ld, kernel plumpness, and
29	lodging resistance. Robust has ex	
30	yield by 7% in 28 trials in Minr	nesota and by 10% in 49
31	regional trials. It is 2 to 3 days late	r in maturity than Morex.
32	Robust has high levels of resista	nce to stem rust, incited
33	by Puccinia graminis Pers. f. sp. tri	
34	incited by Bipolaris sorokiniana (Sa	cc. in Sorok.) Shoemaker,
35	but is susceptible to loose smut,	caused by Ustilago nuda
36	(Jens.) Rostr.	
37	In regional tests conducted in	n collaboration with the
38	USDA Barley and Malt Lab., N	fadison, Wi., Robust ap-
39	peared equal to Morex in most	quality traits. It has the
10	high extract level of Morex. It is le	ower than Morex in alpha
11	amylase and diastic power but hig	ther in plump kernel per-
12	centage. In pilot tests and in one	plant scale test made in
13	cooperation with the malting and	I brewing industries, Ro-
14	bust was rated satisfactory and eq	ual to the check for malt-
5	ing and brewing characteristics. B	
6	by the Minnesota Agric. Exxp. S	tn., St. Paul, MN 55108.
7		
8	D. C. RASMUSSON	AND R. W. WILCOXSON
9	References and	Notes
0	1 Pasmusson D.C. and D.D. Will.	070 P
2	 Rasmusson, D.C., and R.D. Wilcoxson 1 ley. Crop Sci. 19:293. 	979. Registration of Morex bar-
3	2. Professors, Dep. of Agronomy and Pla	ant Genetics and Dep. of Plant
4	Pathology, respectively, Univ. of Minne	sota, St. Paul, MN 55108. Pub-
5 6	lished with the approval of the Directo	
7	Stn. as Journal Article No. 13, 371. Ac	tepted 15 stay 1985.

The interest in a literal point of the constitution of the constit

UNITED STATES DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE
LIVESTOCK, POULTRY, GRAIN & SEED DIVISION
BELTSVILLE, MARYLAND 20705

EXHIBIT C (Barley)

OBJECTIVE DESCRIPTION OF VARIETY

INSTRUCTIONS: See Reverse.	
Minnesota Agricultural Experiment Station	FOR OFFICIAL USE ONLY
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	ZUNAALI
University of Minnesota	VARIETY NAME OR TEMPORARY
St. Paul, MN 55108	DESIGNATION
Place the appropriate number that describes the varietal characte Place a zero in first box (i.e. 089 or 09) when number	r of this variety in the boxes below.
1. GROWTH HABIT:	
1 = SPRING 2 = FACULTATIVE WINTER 3 = WINTER	3 Early Growth: 1 = PROSTRATE 2 = SEMIPROSTRATE 3 = ERECT
2. MATURITY (50% Flowering):	
2 1 = EARLY (California Mariout) 2 = MIDSEASON (Betzes)	3 = LATE (Frontier)
	LIFORNIA MARIOUT 3 = CONQUEST 4 = DICKSON
No. of days Later than 4 5 = PIROLINE 6 = P	RIMUS 7 = UNITAN
3, PLANT HEIGHT (From soil level to top of head):	
3 1 = SEMIDWARF 2 = SHORT (California Mariout) 3 = ME	DIUM TALL (Betzes) 4 = TALL (Conquest)
	ALIFORNIA MARIOUT 3 = CONQUEST 4 = DICKSON PRIMUS 7 = UNITAN
Cm. Taller than	
4. STEM:	
2 Exertion (Flag to spike at maturity): 3 = 10 - 15 cm.	Anthocyanin: 1 = ABSENT 2 = PRESENT
5 NO. OF NODES (Originating from node above ground)	
1 = CLOSED 2 = V-SHAPED 3 = OPEN 4 = MODIFIED CLOSED OR OPEN	1 = STRAIGHT 2 = SNAKY 3 = OTHER (Specify)
5. LEAF:	
Basal leaf sheath (seedling): 1 = GLABROUS 2 = PUBESCENT	Position of flag leaf (at boot stage): 1 = DROOPING 2 = UPRIGHT
3 Waxiness: 1 = ABSENT (Glossy) 2 = SLIGHTLY WAXY 3 = WAXY	1 8 MM, WIDTH (First leaf below flag leaf)
2 4 CM. LENGTH (First leaf below flag leaf)	Anthocyanin in leaf sheath: 1 = ABSENT 2 = PRESENT
6. HEAD:	
2 Type: 1 = TWO-ROWED 2 = SIX-ROWED	2 Density: 2 = ERECT (Not dense) 3 = ERECT (Dense)
Shape: 1 = TAPERING 2 = STRAP 3 = CLAVATE 4 = OTHER (Specify)	2 Waxiness: 1 = ABSENT (Glossy) 2 = SLIGHTLY WAXY 3 = WAXY
3 Lateral Kernels Overlap: 1 = NONE 2 = AT TIP 3 = 1/4 - 1/2 OF HEAD	Rachis (Hair on edge): 1 = LACKING 2 = FEW 3 = COVERED
7. GLUME:	
2 Length: 1 = 1/3 OF LEMMA 2 = 1/2 OF LEMMA 3 = MORE THAN 1/2 OF LEMMA	Hairs: 1 = NONE 2 = SHORT 3 = LONG
4 Hair covering: 1 = NONE 2 = RESTRICTED TO MIDDLE	3 = CONFINED TO BAND 4 = COMPLETELY COVERED
3 Awns: 1 = LESS THAN EQUAL TO LENGTH OF GLUMES 3 = MORE THAN EQUAL TO LENGTH OF GLUMES	2 = EQUAL TO LENGTH OF GLUMES
Awn Surface: 1 = SMOOTH 2 = SEMISMOOTH 3 = RO	UGH

3 = SHORT ON CENTRAL ROWS, AWNLETS ON L 5 = LONG (longer than spike) 6 = HOODED	OWS, AWNLESS ON LATERAL ROWS ATERAL ROWS 4 = SHORT (less than equal to length of spike)				
Awn Surface: 0 = AWNLESS 1 = SMOOTH 2 = SEMISE					
	Awn Surface: 0 = AWNLESS 1 = SMOOTH 2 = SEMISMOOTH 3 = ROUGH				
2 Teeth: 1 = ABSENT 2 = FEW 3 = NUMEROUS	Hair: 1 = ABSENT 2 = PRESENT				
3 Shape of base: 1 = DEPRESSION 2 = SLIGHT CREASE 3 = TRANSVERSE CREASE	Rachilla Hairs: 1 = SHORT 2 = LONG				
9. STIGMA:					
] Hairs: 1 = FEW 2 = MANY					
10. SEED:					
2 Type: 1 = NAKED 2 = COVERED	Hairs on Ventral Furrow: 1 = ABSENT 2 = PRESENT				
3 Length: 1 = SHORT (8.0 mm.) 2 = SHORT TO MIDLONG 4 = MIDLONG TO LONG (9.0 - 10.5 mm.)	3 = MIDLONG (8.5 - 9.5 mm.) 5 = LONG (10.0 mm.)				
2 Wrinkling of hull: 1 = NAKED 2 = SLIGHTLY WRINKLE	ED 3 = SEMIWRINKLED 4 = WRINKLED				
Aleurone Color: 1 = COLORLESS (White or Yellow) 2 =	BLUE				
0 0 PERCENT ABORTIVE	3 5 GMS. PER 1000 SEEDS				
11. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant)					
0 SEPTORIA 0 NET BLOTCH	2 SPOT BLOTCH 1 POWDERY MILDEW				
1 LOOSE SMUT 0 BACTERIAL BLIGHT	0 COVERED SMUT 0 FALSE LOOSE SMUT				
2 STEM RUST 0 LEAF RUST	0 SCALD				
O AY O BSMV	O BYDV OTHER (Specify)				
12. INSECT: (0 = Not tested, 1 = Susceptible, 2 = Resistant)					
0 GREEN BUG 0 ENGLISH GRAIN APHID	0 CHINCH BUG 0 ARMYWORM				
0 GRASS HOPPERS 0 CERIAL LEAF BETTLE	0 OTHER (Specify)				
HESSIAN FLY RACES	0 B 0 C				
0 D 0 E	0 F 0 G				
13. CHEMICAL (0 = Not Tested, 1 = Susceptible, 2 = Resistant)					
0 DDT 0 OTHER (Specify)					
14. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THA	AT SUBMITTED:				
CHARACTER NAME OF VARIETY	CHARACTER NAME OF VARIETY				
Plant tillering MORE X	Seed size LARKER				
Leaf size MANKER	Coleoptile elongation				
Leaf color • MOREX	Seedling pigmentation				
HUILA	grand programme and programme				
Leaf carriage MOREX					

REFERENCES: The following publications may be used as a reference aid for the standardization of character descriptions and terms used in this form:

- 1. Wiebe, G. A., and D. A. Reid, 1961, Classification of Barley Varieties Grown in the United States and Canada in 1958, Technical Bulletin No. 1224, U.S. Dept. of Agriculture.
- 2. Reid, D. A., and G. A. Wiebe, 1968, Barley: Origin, Botany, Culture, Winter Hardiness, Genetics, Utilization, Pests, Agriculture Handbook No. 338, U.S. Dept. of Agriculture. pp. 61 84.
- 3. Malting Barley Improvement Association, Milwaukee, Wisconsin, 1971, Barley Variety Dictionary.

COLOR: Nickerson's or any recognized color fan may be used to determine color of the described variety.

FORM LPGS-470-5 (8-80) (REVERSE)